

付表7-1 基点別、深度別水草採集重量(g)2002年

基点	1						3					
	9/27	02/10/31					9/27	02/10/31				
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7
深度(m)												
被植率(%)	0	0	0	40	0	0	0	0	40	90	100	50
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤									3.3	116.5	1.9
	風乾									0.4	13.0	0.2
ヒロハノエビモ	湿潤								38.9	99.3	3.8	
	風乾								3.6	9.2	0.4	
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤			66.7						1.8	2.9	
	風乾			6.1						0.2	0.3	
イバラモ	湿潤										1.1	101.8
	風乾										0.0	4.0
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤											
	風乾											
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤									0.4	13.5	1.6
	風乾									0.0	0.9	0.1
コカナダモ	湿潤			4.5								
	風乾			0.3								
オオカナダモ	湿潤			35.6								
	風乾			2.0								
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤											
	風乾											
ホザキノフサモ	湿潤									12.6	18.7	
	風乾									0.9	1.4	
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤											
	風乾											

付表7-2 基点別、深度別水草採集重量(g)2002年

基点		5						7					
調査年月日		9/27	02/10/31						02/10/31				
深度(m)		1	2	3	4	5	-	1	2	3	4	5	7
被植率(%)		100	20	100	100	100	-	-	0	30	70	70	100
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤				58.5								
	風乾				6.6								
エビモ	湿潤	16.4											
	風乾	1.4											
センニンモ	湿潤	0.8		11.1	22.9	8.9					1.1	3.7	17.6
	風乾	0.1		1.2	2.6	1.0					0.1	0.4	2.0
ヒロハノエビモ	湿潤	1.4								9.0	58.4	56.8	
	風乾	0.1								0.8	5.4	5.3	
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	0.7	23.3	196.8	13.6	13.2				12.9			
	風乾	0.1	2.1	17.9	1.2	1.2				1.2			
イバラモ	湿潤	2.9		6.2	16.2	13.1					6.3	53.8	15.2
	風乾	0.1		0.2	0.6	0.5					0.2	2.1	0.6
コウガイモ	湿潤		1.3		2.0								
	風乾		0.1		0.1								
ネジレモ	湿潤	36.2	4.3	36.7	10.4								
	風乾	2.2	0.3	2.2	0.6								
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	0.9	0.1	49.4	9.0	76.0						0.7	320.1
	風乾	0.1	0.01	3.1	0.6	4.8						0.04	20.4
コカナダモ	湿潤			2.1	2.6								0.4
	風乾			0.1	0.2								0.03
オオカナダモ	湿潤												
	風乾												
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤										17.3	24.9	2.0
	風乾										1.3	1.8	0.1
不明	湿潤				0.2								
	風乾				0.02								
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-3 基点別、深度別水草採集重量(g)2002年

基点		8						10					
調査年月日		9/27	02/11/7					9/27	02/11/7				
深度(m)		1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)		50	10	90	80	30	0	0	30	60	90	70	0
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤	0.1	48.3	87.3	41.9	36.7		64.4	57.2	54.4	357.9		
	風乾	0.01	5.4	9.7	4.7	4.1		7.2	6.4	6.1	39.9		
ヒロハノエビモ	湿潤												
	風乾												
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤												
	風乾												
イバラモ	湿潤										3.1	0.4	
	風乾										0.1	0.02	
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤												
	風乾												
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	46.5	12.8	15.4	37.9	0.5			7.3	33.5	33.6		
	風乾	3.0	0.8	1.0	2.4	0.03			0.5	2.1	2.1		
コカナダモ	湿潤		0.2	0.3									
	風乾		0.01	0.02									
オオカナダモ	湿潤					4.3					13.1		
	風乾					0.2					0.8		
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤									65.7	27.9		
	風乾									4.8	2.1		
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-4 基点別、深度別水草採集重量(g)2002年

基点	12						14					
	9/27	02/11/7					9/27	02/11/8				
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7
深度(m)												
被植率(%)	0	100	100	100	100	0	100	100	100	100	100	-
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤	27.7	0.2	24.2	3.3		242.8	137.6	142.8	11.7	236.8	
	風乾	3.1	0.02	2.7	0.4		27.1	15.3	15.9	1.3	26.4	
ヒロハノエビモ	湿潤											
	風乾											
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤	527.5	2459.2									
	風乾	48.0	223.6									
イバラモ	湿潤			2.8								
	風乾			0.1								
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤						5.9	21.2				
	風乾						0.4	1.3				
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	4.2	1.7	191.1	306.1			17.4	6.0		161.4	
	風乾	0.3	0.1	12.2	19.5			1.1	0.4		10.3	
コカナダモ	湿潤	5.6					2.9		4.6	1.1	0.8	
	風乾	0.4					0.2		0.3	0.1	0.1	
オオカナダモ	湿潤						21.7	37.6	19.5	284.2	1.6	
	風乾						1.2	2.2	1.1	16.3	0.1	
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤						9.4	6.3	73.4	58.7	108.9	
	風乾						0.5	0.3	4.0	3.2	6.0	
ホザキノフサモ	湿潤	0.2		132.6			2.8	0.4	2.9		281.7	
	風乾	0.01		9.8			0.2	0.0	0.2		20.8	
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤						多い	あり				
	風乾											

付表7-5 基点別、深度別水草採集重量(g)2002年

基点	16						18					
	9/27	02/11/11					10/2	02/11/8				
調査年月日												
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)	100	80	100	20	—	—	100					—
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤	115.2					4.1					
	風乾	13.0					0.5					
エビモ	湿潤											
	風乾											
センニンモ	湿潤	64.4	74.9	0.8			56.1	0.3		0.1		
	風乾	7.2	8.4	0.1			6.3	0.03		0.01		
ヒロハノエビモ	湿潤											
	風乾											
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤											
	風乾											
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤	196.8	6.0									
	風乾	11.7	0.4									
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤		0.4	3.5			9.6		17.4			
	風乾		0.03	0.2			0.6		1.1			
コカナダモ	湿潤						0.7					
	風乾						0.05					
オオカナダモ	湿潤	55.4	266.5	59.9			52.6	589.3	670.4	13.9	566.6	
	風乾	3.2	15.3	3.4			3.0	33.7	38.4	0.8	32.4	
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤	2.4	15.7	0.4			1.1	30.8	109.6			
	風乾	0.1	0.9	0.02			0.1	1.7	6.0			
ホザキノフサモ	湿潤							1.8	4.3	18.2		
	風乾							0.1	0.3	1.3		
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤	多い	あり					あり	あり	あり	あり	
	風乾											

付表7-6 基点別、深度別水草採集重量(g)2002年

基点		20						22					
調査年月日		02/11/11						10/2	02/11/11				
深度(m)		1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)		—	100	100	90	—	—	100	70	50	100	100	—
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤								56.9	1.0			
	風乾								6.4	0.1			
エビモ	湿潤												
	風乾												
センニンモ	湿潤		2.3	270.9	96.2			69.7		6.2	86.0		
	風乾		0.3	30.2	10.7			7.8		0.7	9.6		
ヒロハノエビモ	湿潤												
	風乾												
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤												
	風乾												
イバラモ	湿潤												
	風乾												
コウガイモ	湿潤								210.2				
	風乾								9.5				
ネジレモ	湿潤							280.4					
	風乾							16.7					
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤		5.1		146.8				0.2	1.4	2.1	24.1	
	風乾		0.3		9.3				0.01	0.1	0.1	1.5	
コカナダモ	湿潤											1.0	
	風乾											0.1	
オオカナダモ	湿潤		23.0	22.9	48.4				72.3				
	風乾		1.3	1.3	2.8				4.1				
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤		3.5	20.1	22.4					20.5	165.7	33.6	
	風乾		0.2	1.1	1.2					1.1	9.1	1.8	
ホザキノフサモ	湿潤		11.1							1.5		0.2	
	風乾		0.8							0.1		0.01	
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤			多量						多量		多量	
	風乾												

付表7-7 基点別、深度別水草採集重量(g)2002年

基点	24						26						
	10/2	02/11/7					10/2	02/11/20					
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7	
深度(m)													
被植率(%)	100	100	100	100	100	10	5	80	90	100	100	10	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤							18.9	19.5				
	風乾							1.9	2.0				
ササバモ	湿潤	81.8	8.0	242.6									
	風乾	9.2	0.9	27.4									
エビモ	湿潤												
	風乾												
センニンモ	湿潤	66.2	1.6	1.7	232.9	292.1	3.6	14.4	42.6	80.0	130.9	24.2	
	風乾	7.4	0.2	0.2	26.0	32.6	0.4	1.6	4.8	8.9	14.6	2.7	
ヒロハノエビモ	湿潤	5.8						27.2					
	風乾	0.5						2.5					
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	1.4	51.8	48.7									
	風乾	0.1	4.7	4.4									
イバラモ	湿潤												
	風乾												
コウガイモ	湿潤	7.9		0.8	0.5								
	風乾	0.4		0.04	0.02								
ネジレモ	湿潤		0.5					4.2	15.1		0.1		
	風乾		0.03					0.3	0.9		0.01		
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	85.5	0.3	2.6	14.5	60.1	29.2	7.0	0.8	5.6	3.8	20.4	3.9
	風乾	5.4	0.02	0.2	0.9	3.8	1.9	0.4	0.1	0.4	0.2	1.3	0.2
コカナダモ	湿潤	20.6	0.4	14.1	43.6		2.5				0.9	56.5	7.2
	風乾	1.4	0.03	1.0	2.9		0.2				0.1	3.8	0.5
オオカナダモ	湿潤				18.1		9.1						
	風乾				1.0		0.5						
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤	17.2			1.1	68.7	15.7						0.1
	風乾	0.9			0.1	3.8	0.9						0.01
ホザキノフサモ	湿潤	27.7	0.6	4.1	188.3	0.7		2.0		22.1	67.1	20.3	1.0
	風乾	2.0	0.04	0.3	13.9	0.1		0.1		1.6	5.0	1.5	0.1
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤		あり										
	風乾												

付表7-8 基点別、深度別水草採集重量(g)2002年

基点		28						30					
調査年月日		10/2	02/11/20					10/1	02/11/20				
深度(m)		1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)		100	90	80	80	70	50	100	80	100	90	100	90
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤		126.7	41.8	0.8	51.4	15.8	0.7	2.3	952.7	132.7	291.7	76.3
	風乾		14.1	4.7	0.1	5.7	1.8	0.1	0.3	106.2	14.8	32.5	8.5
ヒロハノエビモ	湿潤			69.1	6.4	0.5	0.8	1.4	2.6	32.2	24.3		
	風乾			6.4	0.6	0.05	0.1	0.1	0.2	3.0	2.3		
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	9.2						48.3	42.2	34.6	2.2		
	風乾	0.8						4.4	3.8	3.1	0.2		
イバラモ	湿潤							2.3					
	風乾							0.1					
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤	85.6	23.4						0.1				
	風乾	5.1	1.4						0.01				
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	2.3	14.6		0.2	2.2	81.3	55.9	0.1	2.5		10.1	4.1
	風乾	0.1	0.9		0.01	0.1	5.2	3.6	0.01	0.2		0.6	0.3
コカナダモ	湿潤			9.1	22.4	60.1	15.3			0.9	27.8	11.0	9.8
	風乾			0.6	1.5	4.1	1.0			0.1	1.9	0.7	0.7
オオカナダモ	湿潤			8.3		< 0.1							
	風乾			0.5									
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤			51.8	86.3	47.6	2.2		83.3	27.7	120.9	106.0	
	風乾			3.8	6.4	3.5	0.2		6.1	2.0	8.9	7.8	
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤		あり										
	風乾												



付表7-9 基点別、深度別水草採集重量(g)2002年

基点	32						34						
	10/1	02/1/19					10/1	02/11/19					
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7	
深度(m)													
被植率(%)	0	80	100	100	100	10	50	100	100	100	100	30	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤			0.8	2.2	0.1					0.1	0.2	
	風乾			0.1	0.2	0.01					0.01	0.02	
ササバモ	湿潤						6.6		1.9				
	風乾						0.7		0.2				
エビモ	湿潤												
	風乾												
センニンモ	湿潤	51.1	59.9	72.6	3.2	4.5	2.8	19.8	52.5	1.0	22.3	25.3	
	風乾	5.7	6.7	8.1	0.4	0.5	0.3	2.2	5.9	0.1	2.5	2.8	
ヒロハノエビモ	湿潤			9.3	1.0		0.3	1.3	22.9	20.8	77.6	0.3	
	風乾			0.9	0.1		0.03	0.1	2.1	1.9	7.2	0.03	
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤			3.4	6.2				16.7	42.0	13.4	1.5	
	風乾			0.3	0.6				1.5	3.8	1.2	0.1	
イバラモ	湿潤			1.3	3.0					1.7		5.1	
	風乾			0.1	0.1					0.1		0.2	
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤								2.1				
	風乾								0.1				
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤		2.8	3.9	11.4	0.8	19.0	9.1	0.9		26.8	52.7	39.6
	風乾		0.2	0.2	0.7	0.1	1.2	0.6	0.1		1.7	3.4	2.5
コカナダモ	湿潤			133.2	399.2	452.1	14.7	0.5		0.5	47.7	187.9	103.1
	風乾			9.0	27.0	30.6	1.0	0.03		0.03	3.2	12.7	7.0
オオカナダモ	湿潤					2.4					12.6	7.7	6.1
	風乾					0.1					0.7	0.4	0.3
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤		28.7	26.7	2.4	15.2		6.9	2.4		2.4		
	風乾		2.1	2.0	0.2	1.1		0.5	0.2		0.2		
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-10 基点別、深度別水草採集重量(g)2002年

基点	36						38					
	10/1	02/11/19					10/1	02/11/18				
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7
深度(m)												
被植率(%)	0	100	100	100	100	10	30	80	80	100	100	100
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤			7.7	0.5							
	風乾			0.8	0.1							
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤	60.0	61.1	171.3	118.2	4.0	10.6	100.1	0.5	0.1		89.9
	風乾	6.7	6.8	19.1	13.2	0.4	1.2	11.2	0.1	0.01		10.0
ヒロハノエビモ	湿潤		19.7									
	風乾		1.8									
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤											
	風乾											
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤	46.2										
	風乾	2.8										
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤		0.6	1.7	2.9	8.1	93.7	75.4				
	風乾		0.04	0.1	0.2	0.5	6.0	4.8				
コカナダモ	湿潤			0.8				2.6	779.4	468.9	731.0	527.3
	風乾			0.1				0.2	52.7	31.7	49.5	35.7
オオカナダモ	湿潤											
	風乾											
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤											
	風乾											
ホザキノフサモ	湿潤		15.1	10.1			3.3	4.2	71.3	0.2	70.4	
	風乾		1.1	0.7			0.2	0.3	5.3	0.01	5.2	
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤											
	風乾											

付表7-11 基点別、深度別水草採集重量(g)2002年

基点	40						42					
	10/1	02/11/18					10/1	02/11/18				
調査年月日	1	2	3	4	5	7	1	2	3	4	5	7
深度(m)												
被植率(%)	50	50	70	100	100	80	0	80	100	100	100	100
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤			2.9		0.3		29.3	77.6	44.9		
	風乾			0.3		0.03		2.9	7.8	4.5		
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤	1.7	0.6	9.6		20.3	4.4	27.2		140.1	924.4	294.2
	風乾	0.2	0.1	1.1		2.3	0.5	3.0		15.6	103.1	32.8
ヒロハノエビモ	湿潤											
	風乾											
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤											
	風乾											
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤											
	風乾											
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	3.9		0.3		0.4	0.5			31.1	0.3	
	風乾	0.2		0.02		0.03	0.03			2.0	0.02	
コカナダモ	湿潤	2.9	404.2	34.5	625.8	45.5	417.9		36.5	27.7	84.4	31.7
	風乾	0.2	27.3	2.3	42.3	3.1	28.3		2.5	1.9	5.7	2.1
オオカナダモ	湿潤											
	風乾											
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤			4.1								1.0
	風乾			0.2								0.1
ホザキノフサモ	湿潤	3.2	6.5	36.4	1.9	8.5	0.8		8.8		3.1	1.3
	風乾	0.2	0.5	2.7	0.1	0.6	0.1		0.6		0.2	0.1
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤											
	風乾											

付表7-12 基点別、深度別水草採集重量(g)2002年

基点		44						46					
調査年月日		9/25	02/11/15					9/25	02/11/15				
深度(m)		1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)		100	50	80	100	100	0	0	80	80	70	100	80
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤								5.6	18.6			
	風乾								0.6	1.9			
ササバモ	湿潤												
	風乾												
エビモ	湿潤	63.9											
	風乾	5.5											
センニンモ	湿潤				73.3	114.9			112.9	112.2	100.9	88.6	165.6
	風乾				8.2	12.8			12.6	12.5	11.3	9.9	18.5
ヒロハノエビモ	湿潤												
	風乾												
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤			39.0									0.1
	風乾			3.5									0.01
イバラモ	湿潤												
	風乾												
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤												
	風乾												
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	4.8			10.1	25.1			110.0	73.5	113.2	206.9	0.1
	風乾	0.3			0.6	1.6			7.0	4.7	7.2	13.2	0.01
コカナダモ	湿潤	30.5		138.1	81.4	40.1				137.8	77.9	16.5	29.2
	風乾	2.1		9.3	5.5	2.7				9.3	5.3	1.1	2.0
オオカナダモ	湿潤												
	風乾												
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤				2.3								
	風乾				0.2								
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-13 基点別、深度別水草採集重量(g)2002年

基点	48						50						
	9/25	02/11/15					9/25	02/11/15					
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	0	70	80	80	100	40	0	80	100	100	10	10	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤		31.5	20.4						17.0			
	風乾		3.6	2.3						1.9			
エビモ	湿潤												
	風乾												
センニンモ	湿潤		106.3	76.2	17.3	3.0	15.1		97.0	23.2	167.1	1.6	
	風乾		11.9	8.5	1.9	0.3	1.7		10.8	2.6	18.6	0.2	
ヒロハノエビモ	湿潤								16.0	30.5	285.7		
	風乾								1.5	2.8	26.5		
サンニンモ	湿潤												
	風乾												
オオササエビモ	湿潤								43.3	8.2			
	風乾								3.9	0.7			
イバラモ	湿潤									13.8			
	風乾									0.5			
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤												
	風乾												
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤						3.2			142.8	0.8		
	風乾						0.2			9.1	0.1		
コカナダモ	湿潤		2.2	20.9	264.3	310.8	174.8			223.7	0.7		
	風乾		0.1	1.4	17.9	21.0	11.8			15.1	0.05		
オオカナダモ	湿潤			7.1						13.9	156.3	97.3	59.6
	風乾			0.4						0.8	8.9	5.6	3.4
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤			1.5	0.4	141.4	5.6			94.8			
	風乾			0.1	0.02	7.8	0.3			5.2			
ホザキノフサモ	湿潤		8.8		15.0					224.9	18.9		
	風乾		0.6		1.1					16.6	1.4		
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-14 基点別、深度別水草採集重量(g)2002年

基点	52						54					
	9/25	02/11/14					9/25	02/11/14				
調査年月日												
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)	100			100	100	90	0	0	20	10	20	0
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤				0.9	12.8						
	風乾				0.1	1.3						
ササバモ	湿潤	122.6										
	風乾	13.9										
エビモ	湿潤											
	風乾											
センニンモ	湿潤	2.4	93.4	20.8	53.7			42.6	3.3	6.8		
	風乾	0.3	10.4	2.3	6.0			4.8	0.4	0.8		
ヒロハノエビモ	湿潤			14.6				8.4	2.6	99.1		
	風乾			1.4				0.8	0.2	9.2		
サンニンモ	湿潤											
	風乾											
オオササエビモ	湿潤	4.4										
	風乾	0.4										
イバラモ	湿潤	241.3				3.2						
	風乾	9.5				0.1						
コウガイモ	湿潤			0.1								
	風乾			0.005								
ネジレモ	湿潤											
	風乾											
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	11.2		6.1	2.9	2.4			1.1	0.8	0.2	
	風乾	0.7		0.4	0.2	0.2			0.1	0.1	0.01	
コカナダモ	湿潤		1.9	19.2		0.3			2.0			
	風乾		0.1	1.3		0.02			0.1			
オオカナダモ	湿潤			1.3								
	風乾			0.1								
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤	0.1	2.6	14.5						0.1		
	風乾	0.01	0.1	0.8						0.01		
ホザキノフサモ	湿潤				10.1						6.5	
	風乾				0.7						0.5	
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤											
	風乾											

付表7-15 基点別、深度別水草採集重量(g)2003年

基点	2						4					
	2003/7/23						2003/7/23					
調査年月日	2003/7/23						2003/7/23					
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)	0	1	70	90	90	10	0	5	30	70	70	0
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤	1.6	111.4	117.4	128.5		3.8	89.7	175.2			
	風乾	0.2	12.4	13.1	14.3		0.4	10.0	19.5			
ヒロハノエビモ	湿潤		26.9		83.6			1.8				
	風乾		2.5		7.8			0.2				
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤			16.5							76.9	
	風乾			1.50							7.0	
イバラモ	湿潤					3.8						
	風乾					0.1						
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤											
	風乾											
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	8.9	19.0	16.6	11.1	18.5	16.0	61.6	6.9	64.4		
	風乾	0.6	1.2	1.1	0.7	1.2	1.0	3.9	0.4	4.1		
コカナダモ	湿潤											
	風乾											
オオカナダモ	湿潤											
	風乾											
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤											
	風乾											
ホザキノフサモ	湿潤		118.9		16.5							
	風乾		8.8		1.2							
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤											
	風乾											

付表7-16 基点別、深度別水草採集重量(g)2003年

基点		6						9					
調査年月日		2003/7/23						03/8/7					
深度(m)		1	2	3	4	5	—	1	2	3	4	5	7
被植率(%)		20	100	80	90	100		5	20	20	20	0	0
シヤジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤				2.3								
	風乾				0.2								
ササバモ	湿潤		195.8										
	風乾		22.1										
エビモ	湿潤												
	風乾												
センニンモ	湿潤		15.3	85.8	70.3	141.7		2.3	25.3	46.9	38.0		
	風乾		1.7	9.6	7.8	15.8		0.3	2.8	5.2	4.2		
ヒロハノエビモ	湿潤		67.3	8.9	1.3			2.4	5.0				
	風乾		6.2	0.8	0.1			0.2	0.5				
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	7.8	6.5	29.6		15.0							
	風乾	0.7	0.6	2.7		1.4							
イバラモ	湿潤		43.5	0.7	5.1	1.1							
	風乾		1.7	0.03	0.2	0.04							
コウガイモ	湿潤		28.3										
	風乾		1.3										
ネジレモ	湿潤			33.9	37.5								
	風乾			2.0	2.2								
ヒメホタルイ	湿潤	8.6											
	風乾												
クロモ	湿潤	0.1	80.2	22.4	66.9	69.1			24.6	23.9	3.2		
	風乾	0.01	5.1	1.4	4.3	4.4			1.6	1.5	0.2		
コカナダモ	湿潤												
	風乾												
オオカナダモ	湿潤								20.4				
	風乾								1.2				
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤				100.7	24.4			2.6				
	風乾				7.4	1.8			0.2				
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												



付表7-17 基点別、深度別水草採集重量(g)2003年

基点		11						13					
調査年月日		03/8/7						03/7/28					
深度(m)		1	2	3	4	5	7	1	2	3	4	5	—
被植率(%)		5	50	70	80	90	1	100	100	100	80	10	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤	0.4	2.0	82.7	120.2	66.8	1.0	57.3	289.6	952.5	732.6	4.6	
	風乾	0.04	0.2	9.2	13.4	7.4	0.1	6.4	32.3	106.2	81.7	0.5	
ヒロハノエビモ	湿潤	32.1											
	風乾	3.0											
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	1.0	65.2	15.4		0.4		1.3					
	風乾	0.1	5.9	1.4		0.04		0.1					
イバラモ	湿潤	1.3				1.8							
	風乾	0.1				0.1							
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤			2.6									
	風乾			0.2									
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	9.9		28.0	60.8	91.3	1.2	47.1	16.1	3.9			
	風乾	0.6		1.8	3.9	5.8	0.1	3.0	1.0	0.2			
コカナダモ	湿潤							66.0	717.8	124.8	371.5	13.0	
	風乾							4.5	48.6	8.4	25.1	0.9	
オオカナダモ	湿潤									13.4			
	風乾									0.8			
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤								120.9	52.3	1.9		
	風乾								6.6	2.9	0.1		
ホザキノフサモ	湿潤								413.1	56.3			
	風乾								30.5	4.2			
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-18 基点別、深度別水草採集重量(g)2003年

基点	15						17					
	03/7/28						03/8/4					
調査年月日												
深度(m)	1	2	3	4	5	—	1	2	3	4	—	—
被植率(%)	60	90	60	5	80		30	80	80	100		
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤			0.2			84.9					
	風乾			0.02			9.6					
エビモ	湿潤											
	風乾											
センニンモ	湿潤	0.5	153.1	10.9		177.7		147.7	69.5			
	風乾	0.1	17.1	1.2		19.8		16.5	7.8			
ヒロハノエビモ	湿潤											
	風乾											
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤		40.7									
	風乾		3.7									
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤	11.8						22.6				
	風乾	0.5						1.0				
ネジレモ	湿潤							1.9				
	風乾							0.1				
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	48.6	489.6	87.2	1.8	88.6		12.0	123.9	39.0		
	風乾	3.1	31.1	5.5	0.1	5.6		0.8	7.9	2.5		
コカナダモ	湿潤	0.1		2.6	0.9	14.5			1.6			
	風乾	0.01		0.2	0.1	1.0			0.1			
オオカナダモ	湿潤	138.2		7.9		50.4			455.2	127.4		
	風乾	7.9		0.5		2.9			26.1	7.3		
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤		6.9		2.6	7.0	2.6	7.3	76.2	26.9		
	風乾		0.4		0.1	0.4	0.1	0.4	4.2	1.5		
ホザキノフサモ	湿潤			200.9								
	風乾			14.8								
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤	若干有	多い				やや多	やや多	なし	若干		
	風乾											

付表7-19 基点別、深度別水草採集重量(g)2003年

基点	19						21					
	03/8/4						03/7/21					
調査年月日												
深度(m)	1	2	3	4	—	—	1	2	3	4	5	—
被植率(%)	0	100	100	60			100	100	100	100	70	
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤						56.0					
	風乾						6.3					
エビモ	湿潤											
	風乾											
センニンモ	湿潤	128.9	197.4	21.5			211.3	189.6	1158.1	440.1	136.6	
	風乾	14.4	22.0	2.4			23.6	21.1	129.2	49.1	15.2	
ヒロハノエビモ	湿潤											
	風乾											
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤											
	風乾											
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤						5.1					
	風乾						0.3					
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	16.5	52.2	21.8			1.2	3.7	1.3	1.1	44.6	
	風乾	1.0	3.3	1.4			0.1	0.2	0.1	0.1	2.8	
コカナダモ	湿潤	1.0	5.5								1.9	
	風乾	0.1	0.4								0.1	
オオカナダモ	湿潤	37.8	780.1	136.9				38.4		48.9	51.9	
	風乾	2.2	44.7	7.8				2.2		2.8	3.0	
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤		53.9				16.2	118.1	310.0	896.1	53.1	
	風乾		3.0				0.9	6.5	17.0	49.2	2.9	
ホザキノフサモ	湿潤			4.3				200.9				
	風乾			0.3				14.8				
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤			微量			多い					
	風乾											

付表7-20 基点別、深度別水草採集重量(g)2003年

基点		23						25					
調査年月日		03/7/29						03/8/7					
深度(m)		1	2	3	4	5	—	1	2	3	4	5	7
被植率(%)		100	90	100	100	90		40	100	70	70	80	40
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤												
	風乾												
ササバモ	湿潤		11.4					16.6		172.5			
	風乾		1.3					1.9		19.5			
エビモ	湿潤												
	風乾												
センニンモ	湿潤	241.6	13.3	421.4	281.4	126.6		9.6	42.1	4.4	203.9	268.2	56.5
	風乾	26.9	1.5	47.0	31.4	14.1		1.1	4.7	0.5	22.7	29.9	6.3
ヒロハノエビモ	湿潤												
	風乾												
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤							0.5	15.0	9.3			
	風乾							0.0	1.4	0.8			
イバラモ	湿潤												
	風乾												
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤							10.9	22.6	20.7			
	風乾							0.6	1.3	1.2			
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	36.3	64.7	0.8	22.0	11.1		309.0	277.8	26.7	47.5	10.2	1.3
	風乾	2.3	4.1	0.1	1.4	0.7		19.7	17.7	1.7	3.0	0.6	0.1
コカナダモ	湿潤	0.7		34.2	25.4	2.1		0.6		1.3	0.9	8.1	57.8
	風乾	0.05		2.3	1.7	0.1		0.04		0.1	0.1	0.5	3.9
オオカナダモ	湿潤				4.5	76.2							1.6
	風乾				0.3	4.4							0.1
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤	2.7	11.0	9.6	440.9	294.6							
	風乾	0.1	0.6	0.5	24.2	16.2							
ホザキノフサモ	湿潤	3.3	450.6	20.1				14.8	90.5	29.3			
	風乾	0.2	33.2	1.5				1.1	6.7	2.2			
不明	湿潤												
	風乾												
ヒシ	湿潤							346.3					
	風乾							51.2					
糸状藻類	湿潤	若干有	若干有										
	風乾												

付表7-21 基点別、深度別水草採集重量(g)2003年

基点	27						29						
	03/8/20						03/8/21						
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	0	40	40	30	10	40	0	10	40	50	50	20	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤									4.3			
	風乾									0.4			
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤		6.9	0.3	62.1	42.5	114.1			33.2	180.4	19.0	
	風乾		0.8	0.03	6.9	4.7	12.7			3.7	20.1	2.1	
ヒロハノエビモ	湿潤		10.3	17.1					19.6				
	風乾		1.0	1.6					1.8				
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤				37.6				3.3				
	風乾				3.4				0.3				
イバラモ	湿潤								0.5				
	風乾								0.0				
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤												
	風乾												
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤		13.2	16.1	6.5		6.9			39.1	78.2	4.1	52.4
	風乾		0.8	1.0	0.4		0.4			2.5	5.0	0.3	3.3
コカナダモ	湿潤		43.2	47.7	6.7		4.4			3.9	43.9	3.2	
	風乾		2.9	3.2	0.5		0.3			0.3	3.0	0.2	
オオカナダモ	湿潤		14.4										
	風乾		0.8										
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤		12.6	2.1			2.7			103.2	16.8		
	風乾		0.9	0.2			0.2			7.6	1.2		
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-22 基点別、深度別水草採集重量(g)2003年

基点	31						33					
	03/8/5						03/8/5					
調査年月日												
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7
被植率(%)	70	50	60	70	70	90	0	100	80	100	90	100
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤											
	風乾											
ササバモ	湿潤	79.6										
	風乾	9.0										
エビモ	湿潤											
	風乾											
センニンモ	湿潤	10.8	24.5	56.1	105.4	156.2	44.4	41.3	107.8	24.3		
	風乾	1.2	2.7	6.3	11.8	17.4	5.0	4.6	12.0	2.7		
ヒロハノエビモ	湿潤		2.1	39.1				39.1	19.5	23.2		
	風乾		0.2	3.6				3.6	1.8	2.2		
サンネンモ	湿潤											
	風乾											
オオササエビモ	湿潤	94.1							4.4			
	風乾	8.6							0.4			
イバラモ	湿潤							0.8				
	風乾							0.03				
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤	18.3										
	風乾	1.1										
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤	1.1		1.2	27.3			43.9	12.1	10.5		
	風乾	0.1		0.1	1.7			2.8	0.8	0.7		
コカナダモ	湿潤	9.3	52.8	54.3	108.0	127.9	816.5	86.7	211.4	184.0	450.9	333.5
	風乾	0.6	3.6	3.7	7.3	8.7	55.2	5.9	14.3	12.4	30.5	22.6
オオカナダモ	湿潤								25.6			
	風乾								1.5			
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤					1.0			1.9			
	風乾					0.1			0.1			
ホザキノフサモ	湿潤		4.0		88.4			91.1	1.3	9.2		
	風乾		0.3		6.5			6.7	0.1	0.7		
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤	なし	やや多									
	風乾											

付表7-23 基点別、深度別水草採集重量(g)2003年

基点	35						37						
	03/8/21						03/8/25						
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	90	20	80	60	60	5	0	1	10	5	5	1	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤			3.6		0.8							
	風乾			0.4		0.1							
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤	1.7		0.4	11.7	10.7	2.8			35.1	38.0	46.7	2.8
	風乾	0.2		0.04	1.3	1.2	0.3			3.9	4.2	5.2	0.3
ヒロハノエビモ	湿潤		0.9	20.3						0.2		< 0.1	
	風乾		0.1	1.9						0.02			
サンニンモ	湿潤												
	風乾												
オオササエビモ	湿潤			22.9	28.2	7.0							
	風乾			2.1	2.6	0.6							
イバラモ	湿潤			3.6	11.3								
	風乾			0.1	0.4								
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤			249.6	0.8								
	風乾			14.9	0.05								
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	1188.6	7.0	43.9	96.3	175.4	< 0.1			0.1		0.4	
	風乾	75.6	0.4	2.8	6.1	11.2				0.01		0.03	
コカナダモ	湿潤		2.5						0.4	0.5	0.1		
	風乾		0.2						0.03	0.03	0.01		
オオカナダモ	湿潤												
	風乾												
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤	30.4	7.2			< 0.1							
	風乾	1.7	0.4										
ホザキノフサモ	湿潤			1.5	0.9				3.0				
	風乾			0.1	0.1				0.2				
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-24 基点別、深度別水草採集重量(g)2003年

基点	39						41						
	03/7/31						03/7/31						
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	0	80	30	40	70	5	80	80	100	100	100	70	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤		13.9										
	風乾		1.4										
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤		18.3	3.0	0.8	70.5	0.2			89.2	319.7	271.4	198.5
	風乾		2.0	0.3	0.1	7.9	0.02			9.9	35.7	30.3	22.1
ヒロハノエビモ	湿潤												
	風乾												
サンネンモ	湿潤						1.9	31.5	21.4	3.3			22.2
	風乾						0.1	2.3	1.5	0.2			1.6
オオササエビモ	湿潤												
	風乾												
イバラモ	湿潤												
	風乾												
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤							13.0	24.5				
	風乾							0.8	1.5				
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤		3.6	5.4	47.9	20.9				122.9	37.4	19.6	
	風乾		0.2	0.3	3.0	1.3				7.8	2.4	1.2	
コカナダモ	湿潤		246.7	187.8	247.4	172.6	3.4	176.5	94.4	383.2	177.0	0.9	
	風乾		16.7	12.7	16.7	11.7	0.2	11.9	6.4	25.9	12.0	0.1	
オオカナダモ	湿潤												
	風乾												
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤												
	風乾												
ホザキノフサモ	湿潤										4.5		
	風乾										0.3		
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤								若干有				
	風乾												



付表7-25 基点別、深度別水草採集重量(g)2003年

基点	43						45					
	03/8/11						03/8/11					
調査年月日												
深度(m)	1	2	3	4	5	7	1	2	3	4	5	—
被植率(%)	0	60	100	100	80	0	1	40	10	20	10	
シャジクモ	湿潤											
	風乾											
ヒロハノセンニンモ	湿潤		1.1			2.1				4.1		
	風乾		0.1			0.2				0.4		
ササバモ	湿潤											
	風乾											
エビモ	湿潤											
	風乾											
センニンモ	湿潤		6.9	34.6	51.2	7.5		19.0		29.7	1.5	
	風乾		0.8	3.9	5.7	0.8		2.1		3.3	0.2	
ヒロハノエビモ	湿潤			0.2								
	風乾			0.02								
サンネンモ	湿潤		2.3	3.3	5.9	2.2						
	風乾		0.2	0.2	0.4	0.2						
オオササエビモ	湿潤		12.7									
	風乾		1.2									
イバラモ	湿潤											
	風乾											
コウガイモ	湿潤											
	風乾											
ネジレモ	湿潤											
	風乾											
ヒメホタルイ	湿潤											
	風乾											
クロモ	湿潤			1.8	13.6	0.5		1.3	29.9	8.1		
	風乾			0.1	0.9	0.03		0.1	1.9	0.5		
コカナダモ	湿潤		169.5	714.3	416.8	157.9		15.4		736.2	47.6	
	風乾		11.5	48.3	28.2	10.7		1.0		49.8	3.2	
オオカナダモ	湿潤					8.0						
	風乾					0.5						
ハゴロモモ	湿潤											
	風乾											
マツモ	湿潤									2.2		
	風乾									0.1		
ホザキノフサモ	湿潤			4.9		1.2			8.8			
	風乾			0.4		0.1			0.6			
不明	湿潤											
	風乾											
ヒシ	湿潤											
	風乾											
糸状藻類	湿潤							若干				
	風乾											

付表7-26 基点別、深度別水草採集重量(g)2003年

基点	47						49						
	03/8/11						03/8/25						
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	1	100	30	30	30	20	80	100	80	100	80	1	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤		8.7		0.3	1.1		0.8		1.9	13.0		
	風乾		0.9		0.03	0.1		0.1		0.2	1.3		
ササバモ	湿潤												
	風乾												
エビモ	湿潤												
	風乾												
センニンモ	湿潤		136.9		6.3	1.4	96.5	2.2	49.8	17.8	84.5	122.8	
	風乾		15.3		0.7	0.2	10.8	0.2	5.6	2.0	9.4	13.7	
ヒロハノエビモ	湿潤							46.3	23.5	11.0	22.5		
	風乾							4.3	2.2	1.0	2.1		
サンネンモ	湿潤					15.3							
	風乾					1.1							
オオササエビモ	湿潤						95.2	19.0	1.5				
	風乾						8.7	1.7	0.1				
イバラモ	湿潤						1.3	9.5		0.3			
	風乾						0.1	0.4		0.01			
コウガイモ	湿潤												
	風乾												
ネジレモ	湿潤						4.6						
	風乾						0.3						
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤		88.9	41.9	7.1		1.3	301.6	1234.1	111.9	220.6	50.4	1.4
	風乾		5.7	2.7	0.5		0.1	19.2	78.5	7.1	14.0	3.2	0.1
コカナダモ	湿潤	4.3	38.3	191.8	829.8	152.6	35.6		342.5				
	風乾	0.3	2.6	13.0	56.1	10.3	2.4		23.2				
オオカナダモ	湿潤						1.9		9.5				
	風乾						0.1		0.5				
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤				3.5		15.4	151.9	4.0				
	風乾				0.2		0.8	8.3	0.2				
ホザキノフサモ	湿潤												
	風乾												
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-27 基点別、深度別水草採集重量(g)2003年

基点	51						53						
	03/8/25						03/7/24						
調査年月日													
深度(m)	1	2	3	4	5	7	1	2	3	4	5	7	
被植率(%)	30	80	80	70	80	30	0	50	70	70		40	
シャジクモ	湿潤												
	風乾												
ヒロハノセンニンモ	湿潤								0.6				
	風乾								0.1				
ササバモ	湿潤							5.0			強風のため採取できず		
	風乾							0.6					
エビモ	湿潤												
	風乾												
センニンモ	湿潤		2.5	33.7	94.0	277.5	3.4		3.4	1.5			4.5
	風乾		0.3	3.8	10.5	30.9	0.4		0.4	0.2			0.5
ヒロハノエビモ	湿潤		331.0	101.6	180.7	19.8		488.3		46.5			
	風乾		30.7	9.4	16.8	1.8		45.3		4.3			
サンネンモ	湿潤												
	風乾												
オオササエビモ	湿潤	88.1	33.5	0.2					239.2				
	風乾	8.0	3.0	0.02					21.7				
イバラモ	湿潤						2.1			0.1		1.2	
	風乾						0.1			0.00		0.05	
コウガイモ	湿潤			12.8									
	風乾			0.6									
ネジレモ	湿潤												
	風乾												
ヒメホタルイ	湿潤												
	風乾												
クロモ	湿潤	0.3	12.9	39.6	56.2	72.0	136.2		18.5	30.6	136.3	29.3	
	風乾	0.02	0.8	2.5	3.6	4.6	8.7		1.2	1.9	8.7	1.9	
コカナダモ	湿潤		5.3										
	風乾		0.4										
オオカナダモ	湿潤												
	風乾												
ハゴロモモ	湿潤												
	風乾												
マツモ	湿潤	0.3	5.3	21.8		1.6			1.1	6.0	5.4		
	風乾	0.02	0.3	1.2		0.1			0.1	0.3	0.3		
ホザキノフサモ	湿潤			2.3									
	風乾			0.2									
不明	湿潤												
	風乾												
ヒシ	湿潤												
	風乾												
糸状藻類	湿潤												
	風乾												

付表7-28 基点別、深度別水草採集重量(g)2003年

基点		55					
調査年月日		03/7/24					
深度(m)		1	2	3	4	5	7
被植率(%)		0	0	0	5	20	5
シヤジクモ	湿潤						
	風乾						
ヒロハノセンニンモ	湿潤						
	風乾						
ササバモ	湿潤						
	風乾						
エビモ	湿潤						
	風乾						
センニンモ	湿潤						0.7
	風乾						0.1
ヒロハノエビモ	湿潤						
	風乾						
サンネンモ	湿潤						
	風乾						
オオササエビモ	湿潤						
	風乾						
イバラモ	湿潤				6.3		
	風乾				0.2		
コウガイモ	湿潤					19.3	
	風乾					0.9	
ネジレモ	湿潤						
	風乾						
ヒメホタルイ	湿潤						
	風乾						
クロモ	湿潤				8.9	14.6	11.8
	風乾				0.6	0.9	0.8
コカナダモ	湿潤						
	風乾						
オオカナダモ	湿潤						
	風乾						
ハゴロモモ	湿潤						
	風乾						
マツモ	湿潤						
	風乾						
ホザキノフサモ	湿潤				121.3		
	風乾				9.0		
不明	湿潤						
	風乾						
ヒシ	湿潤						
	風乾						
糸状藻類	湿潤						
	風乾						