

原発部位	5年予後 診断年	全症例 (上皮内がんを除く)(初発のみ)					手術(原発巣切除)あり					手術(原発巣切除)なし (姑息・対症療法ありを含む)				
		生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率
全部位	2006	432	387	16	835	98.1%	358	158	12	528	97.7%	74	229	4	307	98.7%
	2007	416	307		723	100.0%	345	125		470	100.0%	71	182		253	100.0%
	2008	437	331	12	780	98.5%	362	113	10	485	97.9%	75	218	2	295	99.3%
	2009	426	353	16	795	98.0%	356	159	11	526	97.9%	70	194	5	269	98.1%
	2010	431	409	13	853	98.5%	371	172	7	550	98.7%	60	237	6	303	98.0%
	2011	500	440	27	967	97.2%	389	170	12	571	97.9%	111	270	15	396	96.2%
	2012	507	432	44	983	95.5%	397	145	22	564	96.1%	110	287	22	419	94.7%
	2013	476	369	62	907	93.2%	397	175	42	614	93.2%	79	194	20	293	93.2%
	2014	511	420	82	1,013	91.9%	424	166	56	646	91.3%	87	254	26	367	92.9%
	2015	513	369	92	974	90.6%	408	115	52	575	91.0%	105	254	40	399	90.0%
2006~2015	4,649	3,817	364	8,830	95.9%	3,807	1,498	224	5,529	95.9%	842	2,319	140	3,301	95.8%	
食道	2006	5	24		29	100.0%	4	13		17	100.0%	1	11		12	100.0%
	2007	4	12		16	100.0%	4	3		7	100.0%		9		9	100.0%
	2008	9	14		23	100.0%	6	5		11	100.0%	3	9		12	100.0%
	2009	4	19		23	100.0%	4	6		10	100.0%		13		13	100.0%
	2010	8	21		29	100.0%	5	9		14	100.0%	3	12		15	100.0%
	2011	9	20		29	100.0%	7	6		13	100.0%	2	14		16	100.0%
	2012	8	14	2	24	91.7%	6	7		13	100.0%	2	7	2	11	81.8%
	2013	6	16	1	23	95.7%	3	10	1	14	92.9%	3	6		9	100.0%
	2014	13	15	2	30	93.3%	9	5		14	100.0%	4	10	2	16	87.5%
	2015	6	13	2	21	90.5%	5	4		9	100.0%	1	9	2	12	83.3%
2006~2015	72	168	7	247	97.2%	53	68	1	122	99.2%	19	100	6	125	95.2%	
胃	2006	89	66	3	158	98.1%	88	41	3	132	97.7%	1	25		26	100.0%
	2007	82	48		130	100.0%	79	29		108	100.0%	3	19		22	100.0%
	2008	75	52	5	132	96.2%	70	28	4	102	96.1%	5	24	1	30	96.7%
	2009	79	51	2	132	98.5%	78	34	2	114	98.2%	1	17		18	100.0%
	2010	79	67	1	147	99.3%	78	47		125	100.0%	1	20	1	22	95.5%
	2011	89	59	4	152	97.4%	85	38	4	127	96.9%	4	21		25	100.0%
	2012	102	64	6	172	96.5%	97	30	4	131	96.9%	5	34	2	41	95.1%
	2013	104	71	11	186	94.1%	102	55	10	167	94.0%	2	16	1	19	94.7%
	2014	85	58	22	165	86.7%	82	35	19	136	86.0%	3	23	3	29	89.7%
	2015	88	63	16	167	90.4%	82	26	13	121	89.3%	6	37	3	46	93.5%
2006~2015	872	599	70	1,541	95.5%	841	363	59	1,263	95.3%	31	236	11	278	96.0%	
大腸 (結腸 + 直腸)	2006	86	41	4	131	96.9%	84	34	4	122	96.7%	2	7	0	9	100.0%
	2007	74	30		104	100.0%	71	25	0	96	100.0%	3	5	0	8	100.0%
	2008	77	25	2	104	98.1%	76	18	2	96	97.9%	1	7	0	8	100.0%
	2009	61	34	2	97	97.9%	60	30	2	92	97.8%	1	4	0	5	100.0%
	2010	75	34	1	110	99.1%	73	32	1	106	99.1%	2	2	0	4	100.0%
	2011	69	39	3	111	97.3%	67	30	3	100	97.0%	2	9	0	11	100.0%
	2012	86	41	5	132	96.2%	82	33	4	119	96.6%	4	8	1	13	92.3%
	2013	82	34	10	126	92.1%	80	27	10	117	91.5%	2	7	0	9	100.0%
	2014	75	43	8	126	93.7%	75	31	7	113	93.8%	0	12	1	13	92.3%
	2015	81	30	16	127	87.4%	80	16	15	111	86.5%	1	14	1	16	93.8%
2006~2015	766	351	51	1,168	95.6%	748	276	48	1,072	95.5%	18	75	3	96	96.9%	
肝 (肝内 胆管 除く)	2006	11	11	1	23	95.7%	7	2		9	100.0%	4	9	1	14	92.9%
	2007	6	22		28	100.0%	4	6		10	100.0%	2	16		18	100.0%
	2008	8	15	1	24	95.8%	7		1	8	87.5%	1	15		16	100.0%
	2009	8	13	1	22	95.5%	7	8		15	100.0%	1	5	1	7	85.7%
	2010	14	21		35	100.0%	11	6		17	100.0%	3	15		18	100.0%
	2011	10	22		32	100.0%	6	8		14	100.0%	4	14		18	100.0%
	2012	14	18	1	33	97.0%	10	9	1	20	95.0%	4	9		13	100.0%
	2013	8	15	1	24	95.8%	5	4	1	10	90.0%	3	11		14	100.0%
	2014	8	14	1	23	95.7%	8	3	1	12	91.7%		11		11	100.0%
	2015	8	13	2	23	91.3%	6	3	2	11	81.8%	2	10		12	100.0%
2006~2015	95	164	8	267	97.0%	71	49	6	126	95.2%	24	115	2	141	98.6%	
膵	2006	1	28		29	100.0%	1	12		13	100.0%		16		16	100.0%
	2007	1	24		25	100.0%	1	8		9	100.0%		16		16	100.0%
	2008	2	37		39	100.0%	2	11		13	100.0%		26		26	100.0%
	2009	6	30	1	37	97.3%	6	13	1	20	95.0%		17		17	100.0%
	2010	2	39	1	42	97.6%	2	12		14	100.0%		27	1	28	96.4%
	2011	4	36		40	100.0%	4	17		21	100.0%		19		19	100.0%
	2012	6	34		40	100.0%	4	7		11	100.0%	2	27		29	100.0%
	2013	1	44		45	100.0%	1	12		13	100.0%		32		32	100.0%
	2014	4	39	2	45	95.6%	4	15	1	20	95.0%		24	1	25	96.0%
	2015	3	30		33	100.0%	3	5		8	100.0%		25		25	100.0%
2006~2015	30	341	4	375	98.9%	28	112	2	142	98.6%	2	229	2	233	99.1%	

※ここで示しているのは、診断から5年経過後の予後の把握状況であり、「5年生存率」ではありません。

原発部位	全症例 (上皮内がんを除く)(初発のみ)	手術(原発巣切除)あり	手術(原発巣切除)なし (姑息・対症療法ありを含む)
全部位	<p>不明 4.1%</p> <p>死亡 43.2%</p> <p>生存 52.7%</p>	<p>不明 4.1%</p> <p>死亡 27.1%</p> <p>生存 68.9%</p>	<p>不明 4.2%</p> <p>死亡 70.3%</p> <p>生存 25.5%</p>
食道	<p>不明 2.8%</p> <p>死亡 68.0%</p> <p>生存 29.1%</p>	<p>不明 0.8%</p> <p>死亡 55.7%</p> <p>生存 43.4%</p>	<p>不明 4.8%</p> <p>死亡 80.0%</p> <p>生存 15.2%</p>
胃	<p>不明 4.5%</p> <p>死亡 38.9%</p> <p>生存 56.6%</p>	<p>不明 4.7%</p> <p>死亡 28.7%</p> <p>生存 66.6%</p>	<p>不明 4.0%</p> <p>死亡 84.9%</p> <p>生存 11.2%</p>
大腸 (結腸 + 直腸)	<p>不明 4.4%</p> <p>死亡 30.1%</p> <p>生存 65.6%</p>	<p>不明 4.5%</p> <p>死亡 25.7%</p> <p>生存 69.8%</p>	<p>不明 3.1%</p> <p>死亡 78.1%</p> <p>生存 18.8%</p>
肝 (肝内胆管除く)	<p>不明 3.0%</p> <p>死亡 61.4%</p> <p>生存 35.6%</p>	<p>不明 4.8%</p> <p>死亡 38.9%</p> <p>生存 56.3%</p>	<p>不明 1.4%</p> <p>死亡 81.6%</p> <p>生存 17.0%</p>
膵	<p>不明 1.1%</p> <p>死亡 90.9%</p> <p>生存 8.0%</p>	<p>不明 1.4%</p> <p>死亡 78.9%</p> <p>生存 19.7%</p>	<p>不明 0.9%</p> <p>死亡 98.3%</p> <p>生存 0.9%</p>

原発部位	5年予後 診断年	全症例 (上皮内がんを除く)(初発のみ)					手術(原発巣切除)あり					手術(原発巣切除)なし (姑息・対症療法ありを含む)				
		生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率
肺	2006	40	93	2	135	98.5%	34	20	2	56	96.4%	6	73		79	100.0%
	2007	33	71		104	100.0%	30	13		43	100.0%	3	58		61	100.0%
	2008	40	89	2	131	98.5%	36	13	1	50	98.0%	4	76	1	81	98.8%
	2009	43	94	1	138	99.3%	40	16	1	57	98.2%	3	78		81	100.0%
	2010	41	101	2	144	98.6%	35	16	1	52	98.1%	6	85	1	92	98.9%
	2011	54	115	4	173	97.7%	42	22	2	66	97.0%	12	93	2	107	98.1%
	2012	50	139	8	197	95.9%	40	16	3	59	94.9%	10	123	5	138	96.4%
	2013	57	94	6	157	96.2%	50	23	2	75	97.3%	7	71	4	82	95.1%
	2014	79	108	9	196	95.4%	66	21	6	93	93.5%	13	87	3	103	97.1%
	2015	59	103	9	171	94.7%	50	24	5	79	93.7%	9	79	4	92	95.7%
2006~2015	496	1,007	43	1,546	97.2%	423	184	23	630	96.3%	73	823	20	916	97.8%	
乳房	2006	45	3	1	49	98.0%	45	3	1	49	98.0%				0	#DIV/0!
	2007	50	5		55	100.0%	50	5		55	100.0%				0	#DIV/0!
	2008	54	3	1	58	98.3%	54	2	1	57	98.2%		1		1	100.0%
	2009	36	7	1	44	97.7%	36	6	1	43	97.7%		1		1	100.0%
	2010	46	4	2	52	96.2%	45	3	2	50	96.0%	1	1		2	100.0%
	2011	45	14	5	64	92.2%	42	11	2	55	96.4%	3	3	3	9	66.7%
	2012	47	9	1	57	98.2%	47	6		53	100.0%		3	1	4	75.0%
	2013	35	3	5	43	88.4%	35	3	4	42	90.5%			1	1	0.0%
	2014	54	8	6	68	91.2%	50	4	1	55	98.2%	4	4	5	13	61.5%
	2015	63	10	8	81	90.1%	58	5	5	68	92.6%	5	5	3	13	76.9%
2006~2015	475	66	30	571	94.7%	462	48	17	527	96.8%	13	18	13	44	70.5%	
子宮頸部	2006	13	4		17	100.0%	11	1		12	100.0%	2	3		5	100.0%
	2007	7	2		9	100.0%	4			4	100.0%	3	2		5	100.0%
	2008	7	3		10	100.0%	5	2		7	100.0%	2	1		3	100.0%
	2009	14	5		19	100.0%	11	2		13	100.0%	3	3		6	100.0%
	2010	13	3		16	100.0%	12			12	100.0%	1	3		4	100.0%
	2011	14	5		19	100.0%	12			12	100.0%	2	5		7	100.0%
	2012	22	6	5	33	84.8%	16	3	4	23	82.6%	6	3		10	90.0%
	2013	20	2	7	29	75.9%	17	2	5	24	79.2%	3		2	5	60.0%
	2014	26	7	3	36	91.7%	23	4	3	30	90.0%	3	3		6	100.0%
	2015	15	5	1	21	95.2%	13	2		15	100.0%	2	3		6	83.3%
2006~2015	151	42	16	209	92.3%	124	16	12	152	92.1%	27	26	4	57	93.0%	
子宮体部	2006	10	4	2	16	87.5%	10	2	1	13	92.3%		2	1	3	66.7%
	2007	20	4		24	100.0%	20	3		23	100.0%		1		1	100.0%
	2008	15	3		18	100.0%	15	2		17	100.0%		1		1	100.0%
	2009	17	4	3	24	87.5%	17	4	3	24	87.5%				0	#DIV/0!
	2010	21	3		24	100.0%	21	2		23	100.0%		1		1	100.0%
	2011	21	3	1	25	96.0%	21	3	1	25	96.0%				0	#DIV/0!
	2012	19	5	1	25	96.0%	19	4	1	24	95.8%		1		1	100.0%
	2013	22	4	1	27	96.3%	22	4	1	27	96.3%				0	#DIV/0!
	2014	21	6	1	28	96.4%	21	2	1	24	95.8%		4		4	100.0%
	2015	23	2	1	26	96.2%	23	1	1	25	96.0%		1		1	100.0%
2006~2015	189	38	10	237	95.8%	189	27	9	225	96.0%	0	11	1	12	91.7%	
前立腺	2006	45	17	1	63	98.4%	13			13	100.0%	32	17	1	50	98.0%
	2007	43	11		54	100.0%	16	1		17	100.0%	27	10		37	100.0%
	2008	61	14		75	100.0%	28	3		31	100.0%	33	11		44	100.0%
	2009	57	8	1	66	98.5%	27	2		29	100.0%	30	6	1	37	97.3%
	2010	58	12	2	72	97.2%	32	3		35	100.0%	26	9	2	37	94.6%
	2011	70	18	5	93	94.6%	29			29	100.0%	41	18	5	64	92.2%
	2012	57	11	6	74	91.9%	22	1		23	100.0%	35	10	6	51	88.2%
	2013	49	9	7	65	89.2%	24			24	100.0%	25	9	7	41	82.9%
	2014	48	9	8	65	87.7%	28	4	3	35	91.4%	20	5	5	30	83.3%
	2015	67	11	18	96	81.3%	31		4	35	88.6%	36	11	14	61	77.0%
2006~2015	555	120	48	723	93.4%	250	14	7	271	97.4%	305	106	41	452	90.9%	

※ここで示しているのは、診断から5年経過後の予後の把握状況であり、「5年生存率」ではありません。

原発部位	全症例 (上皮内がんを除く)(初発のみ)	手術(原発巣切除)あり	手術(原発巣切除)なし (姑息・対症療法ありを含む)
肺	<p>不明 2.8%</p> <p>生存 32.1%</p> <p>死亡 65.1%</p>	<p>不明 3.7%</p> <p>生存 67.1%</p> <p>死亡 29.2%</p>	<p>不明 2.2%</p> <p>生存 8.0%</p> <p>死亡 89.8%</p>
乳房	<p>不明 5.3%</p> <p>生存 83.2%</p> <p>死亡 11.6%</p>	<p>不明 3.2%</p> <p>生存 87.7%</p> <p>死亡 9.1%</p>	<p>不明 29.5%</p> <p>生存 29.5%</p> <p>死亡 40.9%</p>
子宮頸部	<p>不明 7.7%</p> <p>生存 72.2%</p> <p>死亡 20.1%</p>	<p>不明 7.9%</p> <p>生存 81.6%</p> <p>死亡 10.5%</p>	<p>不明 7.0%</p> <p>生存 47.4%</p> <p>死亡 45.6%</p>
子宮体部	<p>不明 4.2%</p> <p>生存 79.7%</p> <p>死亡 16.0%</p>	<p>不明 4.0%</p> <p>生存 84.0%</p> <p>死亡 12.0%</p>	<p>不明 8.3%</p> <p>生存 0.0%</p> <p>死亡 91.7%</p>
前立腺	<p>不明 6.6%</p> <p>生存 76.8%</p> <p>死亡 16.6%</p>	<p>不明 2.6%</p> <p>生存 92.3%</p> <p>死亡 5.2%</p>	<p>不明 9.1%</p> <p>生存 67.5%</p> <p>死亡 23.5%</p>