


シロイチモジヨトウ半旬別誘殺数
(安土町大中：P. T)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0	0
	2	1	1	0	0
	3	0	1	0	0
	4		#N/A	1	1
	5		#N/A	0.8	1.4
	6		#N/A	0.8	2.0
5	1		#N/A	1.0	2.8
	2		#N/A	1.0	3.6
	3		#N/A	1.3	4.6
	4		#N/A	2.0	6.2
	5		#N/A	3.0	8.6
	6		#N/A	4.0	11.8
6	1		#N/A	4.3	15.2
	2		#N/A	5.8	19.8
	3		#N/A	4	23.0
	4		#N/A	4	25.8
	5		#N/A	1	26.8
	6		#N/A	3	28.8
7	1		#N/A	7.8	35.0
	2		#N/A	9.8	42.8
	3		#N/A	11.5	52.0
	4		#N/A	8.8	59.0
	5		#N/A	10.5	67.4
	6		#N/A	8.3	74.0
8	1		#N/A	11.8	83.4
	2		#N/A	21.3	100.4
	3		#N/A	20.0	116.4
	4		#N/A	21.8	133.8
	5		#N/A	26.0	154.6
	6		#N/A	41.0	187.4
9	1		#N/A	20.3	203.6
	2		#N/A	12.5	213.6
	3		#N/A	15.8	229.4
	4		#N/A	14.2	243.6
	5		#N/A	45.2	288.8
	6		#N/A	43.4	332.2
10	1		#N/A	68.0	400.2
	2		#N/A	49.6	449.8
	3		#N/A	37.8	487.6
	4		#N/A	15.4	503.0
	5		#N/A	12.4	515.4
	6		#N/A	9.6	525.0
11	1		#N/A	3.8	528.8
	2		#N/A	8.0	536.8
	3		#N/A	1.8	538.6
	4		#N/A	2.6	541.2
	5		#N/A	1.8	543.0
	6		#N/A	1.6	544.6



・平年値はR1～R6の6年間の平均値。
ただし、R2の4月第1半旬～9月第2半旬
は欠測値。

ヨトウガ半旬別誘殺数
(安土町大中：P. T)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0.4	0.4
	2	0	0	0.5	0.9
	3	0	0	0.3	1.2
	4			0.3	1.5
	5			0.6	2.1
	6			0.9	3.0
5	1			0.6	3.6
	2			0.3	3.9
	3			0.2	4.1
	4			0.3	4.4
	5			0.2	4.6
	6			0.1	4.7
6	1			0.0	4.7
	2			0.2	4.9
	3			0.3	5.2
	4			0	5.2
	5			0.1	5.3
	6			0.1	5.4
7	1			0.1	5.5
	2			0	5.5
	3			0	5.5
	4			0	5.5
	5			0.1	5.6
	6			0.1	5.7
8	1			0.0	5.7
	2			0	5.7
	3			0	5.7
	4			0	5.7
	5			0	5.7
	6			0.2	5.9
9	1			0	6.0
	2			0.3	6.3
	3			0.5	6.8
	4			0.2	7.0
	5			0.3	7.3
	6			0.7	8.0
10	1			0.4	8.4
	2			0.4	8.8
	3			0.3	9.1
	4			0.1	9.2
	5			0.1	9.3
	6			0.0	9.3
11	1			0.2	9.5
	2			0.3	9.8
	3			0	9.9
	4			0	9.9
	5			0	9.9
	6			0	9.9



・ 平年値はH27～R 6 の10年間の平均値。

オオタバコガ半旬別誘殺数
(安土町大中：P. T)




月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	4.1	0	0.1	0.1
	2	4.2	0	0.2	0.3
	3	4.3	2	0.4	0.7
	4	4.4		0.5	1.2
	5	4.5		0.2	1.4
	6	4.6		0.8	2.2
5	1	5.1		1.9	4.1
	2	5.2		2.7	6.8
	3	5.3		3.6	10.4
	4	5.4		3.9	14.3
	5	5.5		3.7	18.0
	6	5.6		4.1	22.1
6	1	6.1		3.4	25.5
	2	6.2		3.1	28.6
	3	6.3		2.1	30.7
	4	6.4		4.6	35.3
	5	6.5		7.1	42.4
	6	6.6		13.4	55.8
7	1	7.1		14.9	70.7
	2	7.2		11.9	82.6
	3	7.3		10.9	93.5
	4	7.4		8.4	101.9
	5	7.5		7.3	109.2
	6	7.6		10.6	119.8
8	1	8.1		11.7	131.5
	2	8.2		15.0	146.5
	3	8.3		11.3	157.8
	4	8.4		8.8	166.6
	5	8.5		10.1	176.7
	6	8.6		20.0	196.7
9	1	9.1		28.2	224.9
	2	9.2		49.8	274.7
	3	9.3		53.7	328.4
	4	9.4		48.2	376.6
	5	9.5		43.0	419.6
	6	9.6		36.6	456.2
10	1	10.1		57.2	513.4
	2	10.2		53.7	567.1
	3	10.3		60.7	627.8
	4	10.4		54.3	682.1
	5	10.5		58.5	740.6
	6	10.6		68.9	809.5
11	1	11.1		52.5	862.0
	2	11.2		37.8	899.8
	3	11.3		30.7	930.5
	4	11.4		18.8	949.3
	5	11.5		9.5	958.8
	6	11.6		5.7	964.5



・ 平年値はH27～R6 の10年間の平均値。

ニカメイガ半旬別誘殺数（フェロモントラップ）

		近江八幡市安土町大中				長浜市難波町				高島市今津町日置前				(頭)
月	半旬	本年	累積	平年	累積	本年	累積	平年	累積	本年	累積	平年	累積	
4	1	0	0	0.1	0.1									越冬世代
	2	0	0	0	0.2									
	3	0	0	0.3	0.5									
	4			1.8	2.3									
	5			5.1	7.4									
	6			7.4	14.8									
5	1			15.6	30.4			3.8	3.8			10.1	10.1	
	2			18.4	48.8			5.7	9.4			12.0	22.1	
	3			39.2	88.0			10.9	20.3			23.2	45.3	
	4			63.2	151.2			15.9	36.2			27.6	72.9	
	5			75.9	227.1			34.1	70.3			31.0	103.9	
	6			72.5	299.6			26.9	97.2			32.0	135.9	
6	1			33.3	332.9			11.2	108.4			17.1	153.0	第1世代
	2			18.2	351.1			4.2	112.6			5.1	158.1	
	3			7.7	358.8			1.8	114.4			2.3	160.4	
	4			2.2	361.0			0.9	115.3			0.6	161.0	
	5			1.6	362.6			0.8	116.1			0.9	161.9	
	6			2.3	364.9			0.9	117.0			1.0	162.9	
7	1			0.6	365.5			2.0	119.0			2.1	165.0	
	2	#N/A		2.9	2.9	#N/A		3.6	3.6	#N/A		2.2	2.2	
	3	#N/A		12.9	15.8	#N/A		2.9	6.5	#N/A		2.1	4.3	
	4	#N/A		16.3	32.1	#N/A		1.9	8.4	#N/A		2.6	6.9	
	5	#N/A		19.0	51.1	#N/A		1.3	9.7	#N/A		4.0	10.9	
	6	#N/A		15.5	66.6	#N/A		3.4	13.1	#N/A		4.5	15.4	
8	1	#N/A		14.2	80.8	#N/A		5.7	18.8	#N/A		2.6	18.0	
	2	#N/A		10.7	91.5	#N/A		7.8	26.6	#N/A		3.6	21.6	
	3	#N/A		5.5	97.0	#N/A		3.8	30.4	#N/A		4.1	25.7	
	4	#N/A		3.9	100.9	#N/A		2.4	32.8	#N/A		3.3	29.0	
	5	#N/A		7.5	108.4	#N/A		2.6	35.4	#N/A		3.5	32.5	
	6	#N/A		14.7	123.1	#N/A		2.2	37.6	#N/A		6.5	39.0	
9	1	#N/A		12.2	135.3									
	2	#N/A		8.0	143.3									
	3	#N/A		3.6	146.9									
	4	#N/A		0.9	147.8									
	5	#N/A		0.5	148.3									
	6	#N/A		0.2	148.5									




・表中の累積誘殺頭数は、越冬世代と第1世代を区別するため、4月第1半旬～7月第1半旬を越冬世代、7月第2半旬～調査終了時を第1世代として積算している。

・平年値はH27～R6の10年間の平均値。

タバコガ半旬別誘殺数
(安土町大中：P. T)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0	0
	2	0	0	0	0
	3	0	0	0.1	0.1
	4			0.1	0.2
	5			0.3	0.5
	6			0.3	0.8
5	1			0.3	1.1
	2			1.9	3.0
	3			4.0	7.0
	4			5.7	12.7
	5			6.5	19.2
	6			12.5	31.7
6	1			9.3	41.0
	2			8.0	49.0
	3			10.8	59.8
	4			11.9	71.7
	5			7.3	79.0
	6			7.7	86.7
7	1			5.6	92.3
	2			5.6	97.9
	3			8.2	106.1
	4			16.5	122.6
	5			18.7	141.3
	6			25.4	166.7
8	1			17.5	184.2
	2			12.8	197.0
	3			13.3	210.3
	4			14.3	224.6
	5			14.8	239.4
	6			19.1	258.5
9	1			16.0	274.5
	2			23.6	298.1
	3			27.8	325.9
	4			22.6	348.5
	5			22.4	370.9
	6			23.8	394.7
10	1			25.2	419.9
	2			16.5	436.4
	3			13.7	450.1
	4			9.3	459.4
	5			8.2	467.6
	6			6.3	473.9
11	1			5.6	479.5
	2			4.4	483.9
	3			1.9	485.8
	4			1.6	487.4
	5			0.6	488.0
	6			0.3	488.3



・平年値はH27～R6の10年間の平均値。

コナガ半旬別誘殺数
(近江八幡市安土町大中：フェロモントラップ)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0.8	0.8
	2	0	0	0.9	1.7
	3	0	0	1.6	3.3
	4			2.1	5.4
	5			2.8	8.2
	6			2.3	10.5
5	1			2.8	13.3
	2			4.5	17.8
	3			14.0	31.8
	4			18.1	49.9
	5			8.1	58.0
	6			6.2	64.2
6	1			6.0	70.2
	2			4.9	75.1
	3			5.0	80.1
	4			4.4	84.5
	5			4.7	89.2
	6			7.5	96.7
7	1			1.1	97.8
	2			0.6	98.4
	3			0.5	98.9
	4			0.1	99.0
	5			0.4	99.4
	6			1.2	100.6
8	1			0.6	101.2
	2			0.3	101.5
	3			0.2	101.7
	4			0.1	101.8
	5			0.1	101.9
	6			0.2	102.1
9	1			0.1	102.2
	2			0.1	102.3
	3			0.1	102.4
	4			0.2	102.6
	5			0.2	102.8
	6			0.2	103.0
10	1			0.1	103.1
	2			0.1	103.2
	3			0.1	103.3
	4			0.2	103.5
	5			0	103.7
	6			0.4	104.1
11	1			0.5	104.6
	2			0.7	105.3
	3			0.8	106.1
	4			0.8	106.9
	5			0.8	107.7
	6			1.4	109.1



・ 平年値はH27～R 6 の10年間の平均値。

カブラヤガ半旬別誘殺数
(安土町大中：P. T)

月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	10	10	23.4	23.4
	2	9	19	24.5	47.9
	3	15	34	36.5	84.4
	4			40.4	124.8
	5			42.1	166.9
	6			45.3	212.2
5	1			37.3	249.5
	2			30.6	280.1
	3			28.2	308.3
	4			27.4	335.7
	5			20.0	355.7
	6			18.8	374.5
6	1			11.7	386.2
	2			8.0	394.2
	3			9.7	403.9
	4			24.7	428.6
	5			30.1	458.7
	6			28.0	486.7
7	1			24.1	510.8
	2			18.5	529.3
	3			25.7	555.0
	4			20.8	575.8
	5			16.6	592.4
	6			14.7	607.1
8	1			7.8	614.9
	2			7.2	622.1
	3			7.0	629.1
	4			6.2	635.3
	5			7.0	642.3
	6			13.7	656.0
9	1			11.0	667.0
	2			18.5	685.5
	3			24.1	709.6
	4			23.5	733.1
	5			25.8	758.9
	6			24.4	783.3
10	1			24.3	807.6
	2			16.5	824.1
	3			17.8	841.9
	4			13.5	855.4
	5			11.8	867.2
	6			15.0	882.2
11	1			9.9	892.1
	2			10.1	902.2
	3			8.3	910.5
	4			9.7	920.2
	5			8.4	928.6
	6			7.7	936.3






・平年値はH27～R6の10年間の平均値。

ハスモンヨトウ半旬別誘殺数（フェロモントラップ）

（頭）


月	半旬	近江八幡市安土町大中				長浜市難波町				高島市今津町日置前			
		本年	累積	平年	累積	本年	累積	平年	累積	本年	累積	平年	累積
4	1	4.1	0	0.1	0.1								
	2	4.2	0	0.1	0.2								
	3	4.3	0	0	0.2								
	4	4.4		0	0.2								
	5	4.5		0.1	0.3								
	6	4.6		0.1	0.4								
5	1	5.1		0.3	0.7								
	2	5.2		0.6	1.3								
	3	5.3		0.4	1.7								
	4	5.4		0.5	2.2								
	5	5.5		0.6	2.8								
	6	5.6		0.6	3.4								
6	1	6.1		1.5	4.9								
	2	6.2		3.3	8.2								
	3	6.3		5.0	13.2								
	4	6.4		3.7	16.9								
	5	6.5		2.6	19.5								
	6	6.6		3.5	23.0								
7	1	7.1		15.9	38.9		13.2	13.2			7.0	7.0	
	2	7.2		8.3	47.2		17.4	30.6			10.2	17.2	
	3	7.3		9.0	56.2		18.8	49.4			8.5	25.7	
	4	7.4		7.7	63.9		14.0	63.4			4.8	30.5	
	5	7.5		9.7	73.6		10.2	73.6			3.9	34.4	
	6	7.6		10.8	84.4		15.4	89.0			6.2	40.6	
8	1	8.1		11.9	96.3		24.9	113.9			10.9	51.5	
	2	8.2		9.0	105.3		27.4	141.3			15.8	67.3	
	3	8.3		8.7	114.0		19.2	160.5			16.9	84.2	
	4	8.4		9.5	123.5		11.3	171.8			11.3	95.5	
	5	8.5		6.4	129.9		8.7	180.5			16.3	111.8	
	6	8.6		15.5	145.4		16.1	196.6			45.7	157.5	
9	1	9.1		19.1	164.5		41.4	238.0			29	186.4	
	2	9.2		46.8	211.3		28.6	266.6			38.0	224.4	
	3	9.3		38.8	250.1		33.9	300.5			42.6	267.0	
	4	9.4		47.3	297.4		25.0	325.5			46.1	313.1	
	5	9.5		37.2	334.6		17.4	342.9			44.4	357.5	
	6	9.6		33.1	367.7		23.3	366.2			40.6	398.1	
10	1	10.1		65.4	433.1		47.2	413.4			51.8	449.9	
	2	10.2		91.7	524.8		42.5	455.9			66.0	515.9	
	3	10.3		93.8	618.6		31.1	487.0			69.9	585.8	
	4	10.4		75.6	694.2		16.0	503.0			53.0	638.8	
	5	10.5		72.8	767.0		22.8	525.8			40.2	679.0	
	6	10.6		77.2	844.2		31.8	557.6			39.5	718.5	
11	1	11.1		57.7	901.9								
	2	11.2		50.2	952.1								
	3	11.3		32.6	984.7								
	4	11.4		19.8	1004.5								
	5	11.5		15.1	1019.6								
	6	11.6		13.4	1033.0								



・ 平年値はH27～R6の10年間の平均値。

アブラムシ類半旬別誘殺数
(近江八幡市安土町大中：黄色水盤)
(頭)

月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	18	18	29.1	29.1
	2	51	69	19.6	48.7
	3	38	107	31.6	80.3
	4			51.8	132.1
	5			64.4	196.5
	6			82.5	279.0
5	1			94.2	373.2
	2			88.0	461.2
	3			109.8	571.0
	4			94.1	665.1
	5			123.3	788.4
	6			224.9	1013.3
6	1			141.3	1154.6
	2			90.7	1245.3
	3			41.6	1286.9
	4			25.3	1312.2
	5			19.0	1331.2
	6			16.3	1347.5
7	1			8.4	1355.9
	2			6.0	1361.9
	3			9.7	1371.6
	4			9.6	1381.2
	5			10.3	1391.5
	6			14.4	1405.9
8	1			7.2	1413.1
	2			8.4	1421.5
	3			10.1	1431.6
	4			16.9	1448.5
	5			21.9	1470.4
	6			67.6	1538.0
9	1			36.2	1574.2
	2			33.7	1607.9
	3			42.5	1650.4
	4			70.7	1721.1
	5			125.3	1846.4
	6			120.7	1967.1
10	1			118.6	2085.7
	2			49.2	2134.9
	3			33.1	2168.0
	4			37.3	2205.3
	5			39.3	2244.6
	6			59.0	2303.6
11	1			80.4	2384.0
	2			96.5	2480.5
	3			104.6	2585.1
	4			142.8	2727.9
	5			120.6	2848.5
	6			83.7	2932.2



・ 平年値はH27～R 6 の10年間の平均値。