


シロイチモジヨトウ半旬別誘殺数
(安土町大中：P. T)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0	0
	2	1	1	0	0
	3	0	1	0	0
	4	0	1	0	1
	5	0	1	0.6	1.4
	6	0	1	0.6	2.0
5	1	0	1	0.8	2.8
	2	1	2	0.8	3.6
	3	1	3	1.0	4.6
	4	3	6	1.6	6.2
	5	2	8	2.6	8.6
	6	0	8	3.4	11.8
6	1	3	11	3.6	15.2
	2	11	22	4.8	19.8
	3	10	32	4	23.0
	4	5	37	3	25.8
	5	1	38	3	26.8
	6	0	38	5	28.8
7	1	16	54	8.8	35.0
	2	16	70	9.4	42.8
	3	26	96	9.4	52.0
	4	19	115	8.8	59.0
	5	8	123	10.6	67.4
	6	10	133	9.2	74.0
8	1	13	146	10.4	83.4
	2	54	200	17.2	100.4
	3	34	234	16.2	116.4
	4	69	303	21.8	133.8
	5	40	343	23.4	154.6
	6	32	375	44.0	187.4
9	1	49	424	25.4	203.6
	2	122	546	15.0	213.6
	3	142	688	17.0	229.4
	4	142	830	15.2	243.6
	5	335	1165	57.7	288.8
	6	300	1465	54.7	332.2
10	1	285	1750	77.7	400.2
	2	278	2028	56.2	449.8
	3	268	2296	43.8	487.6
	4	196	2492	25.0	503.0
	5	127	2619	22.2	515.4
	6		#N/A	25.8	525.0
11	1		#N/A	7.7	528.8
	2		#N/A	7.2	536.8
	3		#N/A	1.7	538.6
	4		#N/A	2.2	541.2
	5		#N/A	1.5	543.0
	6		#N/A	1.5	544.6



・平年値はR1～R6の6年間の平均値。
ただし、R2の4月第1半旬～9月第2半旬
は欠測値。

ヨトウガ半旬別誘殺数
(安土町大中：P. T)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0.4	0.4
	2	0	0	0.5	0.9
	3	0	0	0.3	1.2
	4	0	0	0.3	1.5
	5	0	0	0.6	2.1
	6	0	0	0.9	3.0
5	1	0	0	0.6	3.6
	2	0	0	0.3	3.9
	3	0	0	0.2	4.1
	4	1	1	0.3	4.4
	5	0	1	0.2	4.6
	6	0	1	0.1	4.7
6	1	0	1	0.0	4.7
	2	0	1	0.2	4.9
	3	0	1	0.3	5.2
	4	0	1	0	5.2
	5	0	1	0.1	5.3
	6	0	1	0.1	5.4
7	1	0	1	0.1	5.5
	2	0	1	0	5.5
	3	0	1	0	5.5
	4	0	1	0	5.5
	5	0	1	0.1	5.6
	6	0	1	0.1	5.7
8	1	0	1	0.0	5.7
	2	0	1	0	5.7
	3	0	1	0	5.7
	4	0	1	0	5.7
	5	0	1	0	5.7
	6	0	1	0.2	5.9
9	1	0	1	0	6.0
	2	0	1	0.3	6.3
	3	0	1	0.5	6.8
	4	0	1	0.2	7.0
	5	0	1	0.3	7.3
	6	0	1	0.7	8.0
10	1	0	1	0.4	8.4
	2	0	1	0.4	8.8
	3	0	1	0.3	9.1
	4	0	1	0.1	9.2
	5	0	1	0.1	9.3
	6			0.0	9.3
11	1			0.2	9.5
	2			0.3	9.8
	3			0	9.9
	4			0	9.9
	5			0	9.9
	6			0	9.9



・平年値はH27～R6の10年間の平均値。

オオタバコガ半旬別誘殺数
(安土町大中：P. T)




月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	4.1	0	0.2	0.1
	2	4.2	0	0.3	0.3
	3	4.3	1	0.3	0.7
	4	4.4	1	0.5	1.2
	5	4.5	1	0.3	1.4
	6	4.6	0	0.9	2.2
5	1	5.1	0	3.2	4.1
	2	5.2	5	4.6	6.8
	3	5.3	6	5.6	10.4
	4	5.4	32	5.2	14.3
	5	5.5	36	4.0	18.0
	6	5.6	33	4.6	22.1
6	1	6.1	7	3.9	25.5
	2	6.2	8	2.8	28.6
	3	6.3	2	2.6	30.7
	4	6.4	1	5.1	35.3
	5	6.5	6	10.0	42.4
	6	6.6	9	16.3	55.8
7	1	7.1	81	16.9	70.7
	2	7.2	50	13.9	82.6
	3	7.3	29	9.9	93.5
	4	7.4	23	8.4	101.9
	5	7.5	13	9.1	109.2
	6	7.6	28	13.7	119.8
8	1	8.1	51	11.6	131.5
	2	8.2	20	14.7	146.5
	3	8.3	19	12.4	157.8
	4	8.4	14	9.5	166.6
	5	8.5	33	12.5	176.7
	6	8.6	39	22.7	196.7
9	1	9.1	97	34.8	224.9
	2	9.2	137	58.0	274.7
	3	9.3	148	60.6	328.4
	4	9.4	76	48.8	376.6
	5	9.5	156	49.6	419.6
	6	9.6	154	47.1	456.2
10	1	10.1	142	62.8	513.4
	2	10.2	110	57.0	567.1
	3	10.3	105	61.0	627.8
	4	10.4	74	58.1	682.1
	5	10.5	153	61.5	740.6
	6	10.6		78.6	809.5
11	1	11.1		61.7	862.0
	2	11.2		44.0	899.8
	3	11.3		36.7	930.5
	4	11.4		22.0	949.3
	5	11.5		11.1	958.8
	6	11.6		6.0	964.5



・ 平年値はH27～R6の10年間の平均値。

ニカメイガ半旬別誘殺数（フェロモントラップ）

近江八幡市安土町大中														長浜市難波町				高島市今津町日置前				(頭)
月	半旬	本年	累積	平年	累積	本年	累積	平年	累積	本年	累積	平年	累積	越冬世代								
4	1	4	0	0	0.1	0.1																
	2	4	0	0	0	0.2																
	3	4	0	0	0.3	0.5																
	4	4	0	0	1.8	2.3																
	5	5	3	3	5.1	7.4																
	6	5	5	8	7.4	14.8																
5	1	5	14	22	15.6	30.4	0	0	3.8	3.8	0	0	10.1		10.1							
	2	5	18	40	18.4	48.8	0	0	5.7	9.4	0	0	12.0		22.1							
	3	5	19	59	39.2	88.0	2	2	10.9	20.3	0	0	23.2		45.3							
	4	5	34	93	63.2	151.2	5	7	15.9	36.2	0	0	27.6	72.9								
	5	6	54	147	75.9	227.1	12	19	34.1	70.3	0	0	31.0	103.9								
	6	6	65	212	72.5	299.6	9	28	26.9	97.2	0	0	32.0	135.9								
6	1	6	20	232	33.3	332.9	3	31	11.2	108.4	0	0	17.1	153.0								
	2	6	18	250	18.2	351.1	1	32	4.2	112.6	0	0	5.1	158.1								
	3	6	8	258	7.7	358.8	1	33	1.8	114.4	1	1	2.3	160.4								
	4	6	3	261	2.2	361.0	1	34	0.9	115.3	1	2	0.6	161.0								
	5	7	1	262	1.6	362.6	0	34	0.8	116.1	2	4	0.9	161.9								
	6	7	0	262	2.3	364.9	0	34	0.9	117.0	2	6	1.0	162.9								
7	1	7	0	262	0.6	365.5	1	35	2.0	119.0	1	7	2.1	165.0								
	2	7	0	0	2.9	2.9	1	1	3.6	3.6	1	1	2.2	2.2								
	3	7	0	0	12.9	15.8	1	2	2.9	6.5	3	4	2.1	4.3								
	4	7	3	3	16.3	32.1	1	3	1.9	8.4	1	5	2.6	6.9								
	5	8	3	6	19.0	51.1	1	4	1.3	9.7	1	6	4.0	10.9								
	6	8	5	11	15.5	66.6	0	4	3.4	13.1	1	7	4.5	15.4								
8	1	8	1	12	14.2	80.8	1	5	5.7	18.8	1	8	2.6	18.0								
	2	8	0	12	10.7	91.5	0	5	7.8	26.6	3	11	3.6	21.6								
	3	8	1	13	5.5	97.0	1	6	3.8	30.4	1	12	4.1	25.7								
	4	8	1	14	3.9	100.9	2	8	2.4	32.8	0	12	3.3	29.0								
	5	9	2	16	7.5	108.4	0	8	2.6	35.4	2	14	3.5	32.5								
	6	9	2	18	14.7	123.1	2	10	2.2	37.6	5	19	6.5	39.0								
9	1	9	2	20	12.2	135.3																
	2	9	0	20	8.0	143.3																
	3	9	0	20	3.6	146.9																
	4	9	0	20	0.9	147.8																
	5	10	1	21	0.5	148.3																
	6	10	0	21	0.2	148.5																
														第1世代								




・表中の累積誘殺頭数は、越冬世代と第1世代を区別するため、4月第1半旬～7月第1半旬を越冬世代、7月第2半旬～調査終了時を第1世代として積算している。

・平年値はH27～R6の10年間の平均値。


タバコガ半旬別誘殺数
(安土町大中：P. T)

月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0	0
	2	0	0	0	0
	3	0	0	0.1	0.1
	4	0	0	0.1	0.2
	5	0	0	0.3	0.5
	6	0	0	0.4	0.9
5	1	2	2	0.8	1.7
	2	5	7	2.9	4.6
	3	9	16	5.4	10.0
	4	8	24	8.6	18.6
	5	2	26	7.8	26.4
	6	4	30	14.5	40.9
6	1	7	37	10.6	51.5
	2	1	38	10.1	61.6
	3	0	38	12.6	74.2
	4	0	38	13.4	87.6
	5	1	39	8.5	96.1
	6	1	40	8.6	104.7
7	1	20	60	6.7	111.4
	2	16	76	8.1	119.5
	3	18	94	9.6	129.1
	4	8	102	16.8	145.9
	5	23	125	21.3	167.2
	6	15	140	28.1	195.3
8	1	11	151	18.2	213.5
	2	8	159	12.3	225.8
	3	3	162	13.6	239.4
	4	1	163	17.0	256.4
	5	6	169	19.2	275.6
	6	7	176	23.6	299.2
9	1	23	199	17.1	316.3
	2	33	232	24.1	340.4
	3	43	275	27.3	367.7
	4	45	320	21.2	388.9
	5	58	378	22.9	411.8
	6	55	433	25.8	437.6
10	1	23	456	30.5	468.1
	2	28	484	20.6	488.7
	3	18	502	16.6	505.3
	4	16	518	12.7	518.0
	5	21	539	8.7	526.7
	6			8.8	535.5
11	1			6.0	541.5
	2			4.6	546.1
	3			2.0	548.1
	4			1.7	549.8
	5			0.6	550.4
	6			0.3	550.7



コナガ半旬別誘殺数
(近江八幡市安土町大中：フェロモントラップ)


月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	0	0	0.8	0.8
	2	0	0	0.9	1.7
	3	0	0	1.6	3.3
	4	5	5	2.1	5.4
	5	1	6	2.8	8.2
	6	5	11	2.3	10.5
5	1	27	38	2.8	13.3
	2	20	58	4.5	17.8
	3	12	70	14.0	31.8
	4	6	76	18.1	49.9
	5	4	80	8.1	58.0
	6	9	89	6.2	64.2
6	1	4	93	6.0	70.2
	2	3	96	4.9	75.1
	3	4	100	5.0	80.1
	4	2	102	4.4	84.5
	5	3	105	4.7	89.2
	6	5	110	7.5	96.7
7	1	3	113	1.1	97.8
	2	1	114	0.6	98.4
	3	2	116	0.5	98.9
	4	3	119	0.1	99.0
	5	1	120	0.4	99.4
	6	0	120	1.2	100.6
8	1	0	120	0.6	101.2
	2	1	121	0.3	101.5
	3	0	121	0.2	101.7
	4	0	121	0.1	101.8
	5	0	121	0.1	101.9
	6	0	121	0.2	102.1
9	1	0	121	0.1	102.2
	2	0	121	0.1	102.3
	3	0	121	0.1	102.4
	4	0	121	0.2	102.6
	5	0	121	0.2	102.8
	6	0	121	0.2	103.0
10	1	0	121	0.1	103.1
	2	1	122	0.1	103.2
	3	0	122	0.1	103.3
	4	0	122	0.2	103.5
	5	0	122	0	103.7
	6			0.4	104.1
11	1			0.5	104.6
	2			0.7	105.3
	3			0.8	106.1
	4			0.8	106.9
	5			0.8	107.7
	6			1.4	109.1



・ 平年値はH27～R 6 の10年間の平均値。

カブラヤガ半旬別誘殺数
(安土町大中：P. T)

月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	10	10	23.1	23.1
	2	9	19	25.4	48.5
	3	15	34	38.2	86.7
	4	11	45	44.5	131.2
	5	6	51	45.3	176.5
	6	10	61	45.2	221.7
5	1	15	76	38.7	260.4
	2	22	98	30.1	290.5
	3	20	118	28.5	319.0
	4	8	126	28.8	347.8
	5	4	130	20.3	368.1
	6	9	139	18.7	386.8
6	1	6	145	11.0	397.8
	2	0	145	7.7	405.5
	3	1	146	10.3	415.8
	4	1	147	25.5	441.3
	5	4	151	29.4	470.7
	6	5	156	27.3	498.0
7	1	4	160	22.6	520.6
	2	17	177	19.8	540.4
	3	6	183	24.7	565.1
	4	17	200	19.4	584.5
	5	10	210	16.1	600.6
	6	6	216	13.7	614.3
8	1	1	217	7.0	621.3
	2	5	222	6.2	627.5
	3	1	223	6.6	634.1
	4	1	224	6.2	640.3
	5	2	226	6.8	647.1
	6	2	228	12.9	660.0
9	1	9	237	9.6	669.6
	2	7	244	15.9	685.5
	3	5	249	21.4	706.9
	4	3	252	20.2	727.1
	5	4	256	22.6	749.7
	6	6	262	22.3	772.0
10	1	4	266	22.4	794.4
	2	6	272	15.1	809.5
	3	7	279	15.9	825.4
	4	1	280	12.2	837.6
	5	11	291	9.8	847.4
	6			13.3	860.7
11	1			8.8	869.5
	2			9.2	878.7
	3			7.7	886.4
	4			10.1	896.5
	5			8.5	905.0
	6			7.3	912.3




・平年値はH27～R6の10年間の平均値。

ハスモンヨトウ半旬別誘殺数（フェロモントラップ）


(頭)

月	半旬	近江八幡市安土町大中				長浜市難波町				高島市今津町日置前			
		本年	累積	平年	累積	本年	累積	平年	累積	本年	累積	平年	累積
4	1	4.1	0	0	0.1	0.1							
	2	4.2	0	0	0.1	0.2							
	3	4.3	0	0	0	0.2							
	4	4.4	0	0	0	0.2							
	5	4.5	0	0	0.1	0.3							
	6	4.6	0	0	0.1	0.4							
5	1	5.1	0	0	0.3	0.7							
	2	5.2	1	1	0.5	1.2							
	3	5.3	2	3	0.3	1.5							
	4	5.4	1	4	0.4	1.9							
	5	5.5	3	7	0.6	2.5							
	6	5.6	19	26	0.6	3.1							
6	1	6.1	5	31	1.5	4.6							
	2	6.2	3	34	3.3	7.9							
	3	6.3	8	42	4.8	12.7							
	4	6.4	10	52	3.1	15.8							
	5	6.5	9	61	2.3	18.1							
	6	6.6	8	69	3.3	21.4							
7	1	7.1	23	92	15.8	37.2	3	3	14.3	14.3	3	3	9.0
	2	7.2	28	120	7.9	45.1	38	41	20.4	34.7	2	5	12.9
	3	7.3	66	186	7.9	53.0	47	88	20.7	55.4	2	7	9.9
	4	7.4	35	221	6.2	59.2	50	138	15.6	71.0	9	16	5.4
	5	7.5	30	251	9.9	69.1	46	184	11.9	82.9	9	25	5.9
	6	7.6	27	278	14.2	83.3	28	212	18.6	101.5	11	36	9.6
8	1	8.1	41	319	14.2	97.5	25	237	22.0	123.5	9	45	11.4
	2	8.2	20	339	10.7	108.2	37	274	25.6	149.1	14	59	21.1
	3	8.3	5	344	11.5	119.7	40	314	18.3	167.4	4	63	22.3
	4	8.4	4	348	11.7	131.4	13	327	14.1	181.5	2	65	15.4
	5	8.5	15	363	6.0	137.4	6	333	9.1	190.6	23	88	29.1
	6	8.6	17	380	11.7	149.1	7	340	15.0	205.6	74	162	64.1
9	1	9.1	21	401	22.9	172.0	12	352	40.7	246.3	51	213	44
	2	9.2	21	422	49.2	221.2	24	376	28.2	274.5	44	257	49.8
	3	9.3	31	453	34.2	255.4	8	384	34.3	308.8	41	298	42.9
	4	9.4	138	591	52.0	307.4	6	390	25.3	334.1	129	427	47.5
	5	9.5	213	804	39.5	346.9	4	394	17.5	351.6	74	501	48.5
	6	9.6	73	877	29.7	376.6	6	400	23.0	374.6	59	560	44.0
10	1	10.1	52	929	61.7	438.3	11	411	47.0	421.6	148	708	31.1
	2	10.2	121	1050	104.7	543.0	15	426	42.6	464.2	69	777	36.5
	3	10.3	107	1157	110.7	653.7	15	441	31.4	495.6	48	825	41.8
	4	10.4	301	1458	105.8	759.5			16.6	512.2	45	870	29.6
	5	10.5	424	1882	95.3	854.8			23.4	535.6	21	891	23.8
	6	10.6			146.1	1000.9			31.8	567.4			31.4
11	1	11.1			116.9	1117.8							
	2	11.2			85.3	1203.1							
	3	11.3			67.8	1270.9							
	4	11.4			46.6	1317.5							
	5	11.5			26.3	1343.8							
	6	11.6			17.4	1361.2							


近江八幡市安土町大中



長浜市難波町



高島市今津町日置前




・平年値はH27～R6の10年間の平均値。

アブラムシ類半旬別誘殺数
(近江八幡市安土町大中：黄色水盤)

(頭)

月	半旬	本年		平年	
		半旬別	累積	半旬別	累積
4	1	4.1	18	29.1	29.1
	2	4.2	51	19.6	48.7
	3	4.3	38	31.6	80.3
	4	4.4	80	51.8	132.1
	5	4.5	54	64.4	196.5
	6	4.6	54	82.5	279.0
5	1	5.1	55	94.2	373.2
	2	5.2	56	88.0	461.2
	3	5.3	95	109.8	571.0
	4	5.4	156	94.1	665.1
	5	5.5	109	123.3	788.4
	6	5.6	79	224.9	1013.3
6	1	6.1	72	141.3	1154.6
	2	6.2	68	985	1245.3
	3	6.3	30	41.6	1286.9
	4	6.4	10	25.3	1312.2
	5	6.5	2	19.0	1331.2
	6	6.6	2	16.3	1347.5
7	1	7.1	7	8.4	1355.9
	2	7.2	4	6.0	1361.9
	3	7.3	4	9.7	1371.6
	4	7.4	3	9.6	1381.2
	5	7.5	3	10.3	1391.5
	6	7.6	3	14.4	1405.9
8	1	8.1	95	7.2	1413.1
	2	8.2	107	8.4	1421.5
	3	8.3	78	10.1	1431.6
	4	8.4	67	16.9	1448.5
	5	8.5	50	21.9	1470.4
	6	8.6	21	67.6	1538.0
9	1	9.1	28	36.2	1574.2
	2	9.2	45	33.7	1607.9
	3	9.3	42	42.5	1650.4
	4	9.4	34	70.7	1721.1
	5	9.5	27	125.3	1846.4
	6	9.6	25	120.7	1967.1
10	1	10.1	21	118.6	2085.7
	2	10.2	17	49.2	2134.9
	3	10.3	33	33.1	2168.0
	4	10.4	22	37.3	2205.3
	5	10.5	16	39.3	2244.6
	6	10.6		59.0	2303.6
11	1	11.1		80.4	2384.0
	2	11.2		96.5	2480.5
	3	11.3		104.6	2585.1
	4	11.4		142.8	2727.9
	5	11.5		120.6	2848.5
	6	11.6		83.7	2932.2



・ 平年値はH27～R 6 の10年間の平均値。

